

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/00, A01K 67/027</b>	<b>A2</b>	(11) International Publication Number: <b>WO 99/05266</b> (43) International Publication Date: 4 February 1999 (04.02.99)
<p>(21) International Application Number: PCT/US98/15387</p> <p>(22) International Filing Date: 24 July 1998 (24.07.98)</p> <p>(30) Priority Data: 60/053,103 26 July 1997 (26.07.97) US</p> <p>(71) Applicant (for all designated States except US): WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, P.O. Box 7365, Madison, WI 53707-7365 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): FIRST, Neal, L. [US/US]; 3157 Highway 78, Mt. Horeb, WI 53572 (US). DOMINKO, Tanja [SL/US]; 2320 White Oak Trail, Oregon, WI 53575 (US). MITALIPOUA, Maissam [US/US]; 2130 University Avenue #6, Madison, WI 53705 (US).</p> <p>(74) Agent: BARTA, Kent, S.; Michael Best &amp; Friedrich LLP, 100 East Wisconsin Avenue, Milwaukee, WI 53202-4108 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> Without international search report and to be republished upon receipt of that report.</p>
<p>(54) Title: TRANS-SPECIES NUCLEAR TRANSFER</p> <p>(57) Abstract</p> <p>A method of producing cloned nuclear transfer embryos from differentiated donor cells is described. The method includes culturing differentiated donor cells in low serum medium and using bovine oocytes as recipient oocytes for donor cells from different species. This method may be used to produce genetically identical animals and transgenic animals.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TRANS-SPECIES NUCLEAR TRANSFERFIELD OF THE INVENTION

5           The present invention relates to the field of the  
cloning of mammals by nuclear transfer, and to a method  
of producing nuclear transfer embryos from a variety of  
cell types, including producing embryos using a donor  
10 cell from one species and a recipient oocyte from  
another species. The method may also be used to  
produce transgenic animals.

BACKGROUND OF THE INVENTION

15           Cloning by nuclear transfer was originally  
described in amphibians (for review see Gurdon, 1986  
and DiBerardino, 1987). More recently, cloning by  
nuclear transfer has been described in the following  
20 mammalian species: sheep (Willadsen, 1986; Campbell et  
al., 1996; and Wilmut et al., 1997); cattle (Robl et  
al., 1986; Prather et al., 1987; Sims and First, 1993;  
U.S. Patents 4,994,384, 5,496,720, 5,057,420,  
5,453,366); rabbits (Stice and Robl, 1988); pigs  
25 (Prather et al., 1989); mice (McGrath and Solter,  
1983); and primates (Meng et al., 1997).

Cloning by nuclear transfer in mammals has several  
common steps. (For review see Prather and First, 1990,  
incorporated herein by reference.) First, a mature  
30 recipient oocyte arrested in Metaphase II is  
enucleated. Then a donor cell is placed under the zona  
pellucida of the recipient oocyte and the two are fused  
together to form a nuclear transfer embryo. Finally,  
it is necessary to activate the nuclear transfer  
35 embryo. Activation is most often accomplished by the  
same electric pulse which was used to fuse the donor  
cell and recipient oocyte, but may be also be  
accomplished by some other stimulus which increases the  
intracellular calcium concentration of the nuclear

-2-

transfer embryo.

For cows and sheep, it has traditionally been necessary to use aged recipient oocytes. The reason for using aged oocytes is that when younger oocytes are used as recipients the nuclear transfer embryo fails to activate. Activation is necessary for the new zygote or nuclear transfer embryo to exit Metaphase II arrest. While sperm are able to activate bovine oocytes after 24 hours of in vitro maturation, oocytes may be activated by stimuli which increase intracellular calcium concentration only after about 42 hours of in vitro culture. (Susko-Parrish et al., 1994). Recently, a method has been described by which bovine oocytes can be successfully activated after only 16 hours of in vitro culture. (Susko-Parish, U.S. Patent 5,496,720, incorporated herein by reference.) This two step method produces an increase in intracellular calcium concentration by treatment with the calcium ionophore ionomycin which is followed by the use of a serine-threonine kinase inhibitor to suppress cellular phosphorylation. It is hypothesized that the serine-threonine kinase inhibitor acts predominantly by inhibiting the c-Mos protein. c-Mos is thought to be necessary to maintain Metaphase II arrest, probably by maintaining the cell cycle protein p34cdc2 in an active state. p34cdc2 in turn is a protein kinase responsible for phosphorylating many proteins which are necessary for oocyte maturation and for normal mitotic cell division. Fertilization of aged oocytes generally results in lower rates of development than when younger oocytes are used. The ability to activate bovine oocytes at an earlier time after the beginning of in vitro culture means nuclear transfer embryos can be activated at a time correlating to the highest rates of development as determined by in vitro fertilization studies.

Traditionally, researchers have found that differentiated cells could not be used for nuclear

-3-

transfer because they would fail to support complete development. Studies in amphibians indicated that as more and more developmentally advanced nuclei are used for nuclear transfer, development decreases. (For review see Gurdon, 1986.) For this reason, many researchers concentrated on culturing embryonic stem cells (ES cells). ES cells are derived from undifferentiated germ cells or embryos and are cultured so as to maintain the cells in an undifferentiated, pluripotential state. ES cells were first described for the mouse and are used to create chimeric embryos since nuclear transfer does not work in mice. (For review see Stewart, 1991; Anderson, 1992.) Nuclear transfer embryos have been produced from ES cells in cattle (Sims and First, 1993) and sheep (Campbell et al., 1996, incorporated herein by reference). However, these methods are somewhat inefficient since only low numbers of the ES cells can be produced.

Recently, researchers have described a method for the production of nuclear transfer sheep embryos from a differentiated adult cell line and from cell lines derived from differentiated embryos (Wilmut et al., 1997, incorporated herein by reference). The donor cells are serum starved by culturing in low serum medium for five days before nuclear transfer. This causes the cells to arrest in G<sub>0</sub>. The researchers hypothesize that nuclei arrested in G<sub>0</sub> are more amenable to nuclear remodeling and reprogramming. Nuclear remodeling and reprogramming describe the events whereby the donor cell nucleus is altered morphologically and biochemically to resemble a normal early embryo nucleus in morphology and transcriptional competence. (For review see Prather and First, 1990; Fulka et al., 1996.) In effect, the donor cell nucleus is de-differentiated.

This new nuclear transfer procedure may have several important impacts. First, genetically identical clones may be made of adult animals with

-4-

economically valuable traits such as high milk production. Second, the large numbers of donor cells which may be generated by normal cell culture techniques make possible the use of standard cell transfection techniques. For example, the donor cells may be transfected with exogenous DNA and a selectable marker. (For review see Joyner, 1991; Stewart, 1991.) Additionally, homologous recombination techniques can be used to add or delete genes from the donor cells at specific sites within the genome. (Koller et al., 1989; Cappecchi, 1989; Stanton et al., 1990, herein incorporated by reference.) After genetic modification of the donor cells by transfection with a selectable marker or homologous recombination, the donor cell can be used in the nuclear transfer procedure to produce a transgenic animal. The availability of these techniques should greatly increase the efficiency of producing transgenic animals other than mice. These animals may be made transgenic for economically valuable traits or for pharmaceutical proteins which may be harvested from the animals milk or blood. Third, this new technique of nuclear transfer may be used to propagate endangered species where it would be impossible to obtain enough gametes to optimize cloning procedures.

#### SUMMARY OF THE INVENTION

It is an object of the present invention to provide an efficient method of producing cloned embryos from differentiated cells by nuclear transfer. It is another object of this invention to provide a method for cloning donor cells of one species by utilizing a recipient oocyte from another species. It is still another object of this invention to provide cloned embryos. The cloned embryos may be implanted into recipient animals to produce offspring or used to originate totipotent or pluripotent cell lines for

-5-

therapeutic uses. It is also an object of the present invention to provide methods for efficient production of transgenic animals and animals in which genes have been "knocked out" by homologous recombination.

5           Therefore, the present invention provides a method of producing nuclear transfer bovine embryos from bovine somatic donor cells and bovine recipient oocytes or from somatic donor cells of one species and recipient oocytes of another species. The donor cells  
10       may be selected from the group consisting of embryonic derived cells, germ cells, somatic cells, and genetically modified cells. In this method, the donor cells are isolated and then cultured in low serum medium to induce the donor cells to undergo G<sub>0</sub> arrest.  
15       The donor cells are then fused to an enucleated recipient oocyte to create a nuclear transfer embryo. When enucleated bovine recipient oocytes are used, the oocytes are preferably selected from the group of bovine oocytes undergoing nuclear maturation within 16  
20       hours of beginning in vitro culture. Fusion may preferably be accomplished by application of an electric pulse 16-24 hours after the beginning of in vitro culture. The nuclear transfer embryo must then be activated. Preferably, activation is accomplished  
25       by elevating intracellular calcium within the embryo by the use of ionomycin and then incubating the embryo with a serine threonine kinase inhibitor such as dimethylaminopurine (DMAP). The activation step is preferably performed about from 4 to 16 hours after  
30       fusion, and most preferably from about 4 to 8 hours after fusion. Alternatively, activation may be performed about 16 to 52 hours after the beginning of in vitro culture, most preferably about 20 to 32 hours after the beginning of in vitro culture.

35           The present invention also provides embryos produced by the above process and totipotent or pluripotent cell lines derived from those embryos. These embryos individually comprise either cytoplasm

-6-

derived from a bovine oocyte, bovine cytoplasm derived  
from a differentiated bovine cell, cell membrane  
derived from a bovine oocyte, cell membrane derived  
from a differentiated bovine cell and nuclei derived  
5 from a differentiated bovine cell or cytoplasm and cell  
membrane from one species and differentiated cytoplasm,  
differentiated cell membrane, and nuclei derived from a  
differentiated cell of another species.

10

DESCRIPTION OF THE PREFERRED EMBODIMENT

Cloning mammals by nuclear transfer is generally  
comprised of the following steps. First, the recipient  
oocyte is enucleated. Second, the donor cell is placed  
15 in contact with the enucleated oocyte and the two are  
fused together to form a nuclear transfer embryo.  
Third, the nuclear transfer embryo is activated so that  
development can occur. The basic cloning steps for  
bovine embryos have been described in U.S. Patents  
20 4,994,384, 5,496,720, 5,057,420, and 5,453,366,  
incorporated herein by reference. Cloning procedures  
for sheep have been described by Willadsen, 1986,  
Campbell et al., 1996, and Wilmut et al., 1997,  
incorporated herein by reference.

25

Recently, researchers in Scotland demonstrated  
that somatic cells and embryonically derived cells  
could be used as donor cells for nuclear transfer.  
(Wilmut et al., 1997; Campbell et al., 1996.) The  
subject matter of these publications also appears in  
30 PCT Publications WO 97/07669, WO 97/07668 and WO  
96/07732, incorporated herein by reference. Prior to  
these results, it was thought impossible to clone  
mammals or fully developmentally competent amphibians  
by nuclear transfer from cells which have undergone  
35 differentiation. (DiBerardino, 1997.) The entire  
concept of the use of embryonic stem cells to produce  
embryos was based on keeping the stem cells in  
undifferentiated state. The use of stem cells for



-7-

cloning purposes is described in PCT Publications WO 95/17500, WO 95/16770 and WO 95/08625, herein incorporated by reference.

5       The use of somatic cells or other differentiated  
cells for nuclear transfer in mammals has many  
potential uses. First, the cells of a genetically  
valuable animal such as a cow with high milk production  
may be used to make genetically identical cows.  
Second, endangered species could be propagated. Third,  
10       transgenic animals could be produced from cell lines  
transfected with exogenous DNA. The high numbers of  
somatic cells available mean that well developed means  
of transfecting cells with exogenous DNA can be used  
such as the use of selectable markers to select  
15       transfected cells and homologous recombination. By  
using these methods researchers should be able to  
produce transgenic animals far more efficiently than by  
currently used techniques such as pronuclear injection.

One of the most important aspects of this  
20       invention is the wide variety of cells which may be  
used. The term "donor cell", as used herein, means any  
cell of any species which is used to provide a nucleus  
to an enucleated oocyte. Types of donor cells include,  
but are not limited to, somatic cells which are cells  
25       derived from adult tissue, embryonic derived cells  
which are differentiated cells harvested from any stage  
of embryo or fetus, germ cells which are cells which  
form or a product of the germline lineage, and  
genetically modified cells which are cells into which  
30       exogenous DNA has been introduced by any transfection  
technique.

The donor cells used in the present invention must  
be serum starved before use. (Wilmut et al., 1997;  
Campbell et al., 1996.) Serum starvation induces the  
35       donor cells to arrest in the G<sub>0</sub> stage of the cell  
cycle. Serum starvation is a simple technique whereby  
the normal concentration of serum in cell culture  
media, generally about 10 percent, is substantially

-8-

lowered for several days. For example, Wilmut et al., cultured the donor cells used in their experiments for 5 days in a medium with 0.5 percent fetal calf serum. The reason why this treatment produces cells which may be used as donor cells for nuclear transfer is unknown. One theory is that the serum starvation causes changes in chromatin structure which allows for reprogramming of the donor nucleus by the recipient oocyte. An additional benefit of serum starvation is that the cells uniformly arrest at a defined stage of the cell cycle,  $G_0$ .  $G_0$  is a stage of quiescence in cell growth the cells enter into just after mitosis. Actively growing cells generally enter into  $G_1$  and prepare for DNA synthesis if not serum starved. By inducing the  $G_0$  arrest, all the cells used for subsequent cloning are at the same stage of the cell cycle, in contrast to actively growing cells which are at various stages of the cell cycle. The advantage is that the  $G_0$  arrested cells have a  $2N$  amount of DNA which corresponds to the normal amount of DNA contributed to a newly fertilized egg by the oocyte and sperm. Therefore, the nuclear transfer embryos begin development with the correct amount of DNA and correct ploidy.

The term "low serum medium" as used herein means cell culture medium that is supplemented with a lower than normal amount of serum, such as fetal calf serum, so that cells cultured in the medium undergo  $G_0$  arrest. " $G_0$  arrest" means that stage of the cell cycle after mitosis and before DNA synthesis where the cell is quiescent and ceases preparation for DNA synthesis and further cell division. The medium used may be any medium in which the donor cell type is normally cultured. Examples of such media include but are not limited to Ham's F 10, TCM-199, and DMEM. In the preferred embodiment of this invention, donor cells are cultured in low serum medium for a period of time sufficient to induce  $G_0$  arrest. In the most preferred embodiment, the donor cells are cultured for five days

-9-

in alpha-MEM with no serum.

The next step in cloning by nuclear transfer involves the culture, selection and enucleation of the recipient oocyte. "Recipient oocyte" means a mature oocyte which is at the correct stage of development to be fertilized by a sperm cell and which after enucleation, serves as the recipient for the donor cell nucleus. For cloning procedures in most mammals, it is necessary to use mature oocytes which are arrested in Metaphase II. (Prather and First, 1990.) This is the stage at which fertilization leading to competent development takes place in most mammalian species. For species such as dogs in which fertilization takes place at a different stage of oocyte development, it would be necessary to utilize an oocyte of the appropriate stage. The use of Metaphase II oocytes also facilitates enucleation because the Metaphase II plate is located adjacent to the first polar body.

The recipient oocyte may be matured either in vitro or in vivo. In the preferred embodiment of the invention, the recipient oocyte is a mature oocyte of any species. In the most preferred embodiment, the recipient oocyte is a mature bovine oocyte. In vivo matured bovine oocytes may be recovered surgically from either superovulated or nonsuperovulated cows or heifers 35 to 48 hours past the onset of estrus or past an injection of human Chorionic Gonadotropin (hCG) or similar hormone. In vitro matured bovine oocytes may be prepared by the methods of Sirard et al., (1988) Parrish et al., (1988) and Rosenkrans and First (1993), incorporated herein by reference. Briefly, ovaries are collected at a slaughterhouse and transported to the laboratory in normal saline at 30 to 34° C. Transport varies from 2 to 6 hours. Oocytes from small follicles of 1 to 6 mm in diameter are aspirated. The immature oocytes are then washed three times in TL HEPES without glucose and to which 0.22 mM pyruvate and 1 mg/ml BSA have been added. The immature oocytes are then placed

-10-

in 50  $\mu$ l drops of maturation medium under paraffin oil. Maturation medium is composed of TC199 containing 10 percent heat treated fetal calf serum, 0.22 mM pyruvate, 0.5  $\mu$ g/ml FSH-P (Scherring-Plough Animal Health Corp. Kenilworth, N.J.) and 1  $\mu$ g/ml estradiol. Ten oocytes are placed in each drop and incubated overnight at 39° C, in high humidity atmosphere comprised of 5% CO<sub>2</sub> and 95% air. The term "beginning of in vitro culture" means the time at which the recipient oocytes are placed in maturation medium.

In a further preferred embodiment, the recipient bovine oocytes are selected from the group of oocytes which mature within 16 hours of the beginning of in vitro culture. To select such oocytes at 16 hours, it is necessary to remove the enclosing cumulus cells from the oocytes so that the extrusion of the first polar body may be observed. Cumulus may be removed by placing the oocytes in TL-HEPES medium supplemented with 2.0 mg/ml of hyaluronidase and then gently aspirating the oocytes with a fire-polished pipette tip. (Sims and First, 1994.) Oocytes which have extruded the first polar body are in Metaphase II arrest and mature. The reason for selecting this subpopulation of oocytes is that they support a higher level of embryo development after fertilization as demonstrated by Dominko and First, 1997.

In the preferred embodiment, recipient oocytes are enucleated at an appropriate time after maturation has occurred, normally 1 to 15 hours after extrusion of the first polar body. In the most preferred embodiment, bovine oocytes are enucleated 16-20 hours after the beginning of in vitro culture. The procedure for enucleation is as follows and is described in Prather et al., U.S. Patent 4,994,384, incorporated herein by reference. Cumulus cells are removed from the oocytes by placing the oocytes in TL-HEPES medium supplemented with 2.0 mg/ml of hyaluronidase and then gently aspirating the oocytes with a fire-polished pipette

-11-

tip. (Sims and First, 1994.) Oocytes to be enucleated are then placed in culture dishes in 100  $\mu$ l drops under oil of TL-HEPES medium supplemented with calcium and magnesium. (See generally Sims and First, 1994.)

5 Micromanipulation may be performed with a Nikon Diaphot microscope equipped Hoffman optics and Narishige micromanipulators. The recipient oocyte is held with a holding pipet with a 120  $\mu$ m outer diameter and 30  $\mu$ m inner diameter. Enucleation is performed by using a  
10 beveled, sharpened enucleation pipet with an outer diameter of approximately 25  $\mu$ m. The inner diameter may be varied according to the size of the donor cell which is used. To enucleate the oocyte, the enucleation pipet is inserted into the oocyte and a  
15 small amount of cytoplasm is removed just below the first polar body. Preferably, enucleation can be verified by staining the mature recipient oocyte with Hoechst 33342, visualizing DNA by excitation emission upon exposure to ultraviolet light for a short time  
20 (i.e. 5 seconds), enucleating the recipient oocyte, and washing the oocyte 2 times in TL-HEPES.

The next steps involve placing the donor cell under the zona pellucida of the recipient oocyte and fusing the recipient oocyte with donor cell. The donor  
25 cell is placed under the zona pellucidae of the recipient oocyte by use of the enucleation pipet. In the preferred embodiment the donor cell and the recipient oocyte may come from the same species or different species and are fused at the time appropriate  
30 for the recipient oocyte species. Examples include but are not limited to a bovine somatic cell with a bovine recipient oocyte, a sheep somatic cell with a bovine recipient oocyte, or a genetically altered or unaltered non-human primate cell or human cell with a bovine  
35 oocyte.

In the most preferred embodiment, a donor cell which has been cultured in low serum medium is fused at 16-24 hours after initiation of in vitro culture to a

-12-

recipient bovine oocyte selected from the group of oocytes maturing within 16 hours from the beginning of culture. Two main advantages exist for utilizing this method of cloning for bovine donor cells and bovine recipient oocytes. First, genetically valuable cows, such as those that exhibit high milk production or other valuable traits, can be multiplied from the various types of donor cells. Second, transgenic animals can be produced in an efficient manner. Standard methods of DNA transfection along with selectable markers can be utilized with large populations of donor cells. (For review see Joyner, 1991; Stewart, 1991.) Also, homologous recombination techniques can be used to introduce exogenous DNA or to knock out endogenous genes. (Koller et al., 1989; Cappecchi, 1989; Stanton et al., 1990.) The exogenous DNA introduced into the donor cells may be a gene encoding a valuable trait or a gene encoding a pharmaceutical protein which will be harvested from the animal.

These advantages are also applicable when the donor cells and recipient oocytes are derived from different species. In another most preferred embodiment, donor cells from species including but not limited to sheep, pigs, rats, non-human primates and humans are fused to the bovine recipient oocyte. An additional advantage of this method is that nuclear transfer clones of these species may be made utilizing the well-defined bovine recipient oocyte. Maturation procedures and timing for fusion and activation have been optimized in the bovine system. When bovine eggs are used, these two parameters do not have to be optimized as they would need to be if a recipient oocyte from another species is utilized. This may be of special importance in the cloning of endangered species where there are too few animals from which gametes can be obtained to conduct experiments to optimize maturation, culture conditions, and timing for

-13-

developmental competence.

Using the bovine oocyte as the recipient oocyte for donor cells of different species raises several interesting questions. Nuclear transfer utilizing donor cells and recipient oocytes from different species provides a powerful experimental system to analyze early embryo development and the common characteristics of early embryo development shared across species. First, it is known that embryos of different species undergo a period of maternally directed cleavage divisions. These cleavage divisions occur in the absence of transcription from the embryonic genome. The time at which the embryo begins transcription is termed the maternal to embryonic transition (MET). (For review see Telford et al., 1993.) For example, bovine embryos begin transcription at the late 4 to early 8 cell stage, murine embryos at the 2 cell stage, rat embryos at the 2 cell stage, sheep embryos at the 8 cell to 16 cell stage and primate embryos at the 4 cell stage. (Telford et al., 1993.) So the timing of MET is specific to each species. Likewise, the complement of proteins which is initially produced by the embryonic genome seems to vary among species where this has been studied. (Crosby et al., 1988; Barnes and First, 1991; Connover et al., 1991.)

Here, the inventors describe the surprising result that the bovine recipient oocyte is capable of supporting the MET of various species when donor cells of various species are fused to the bovine recipient oocyte to produce a nuclear transfer embryo. Factors present in the bovine recipient oocyte cytoplasm are therefore probably responsible for the MET and subsequent transcription and translation of the species characteristic MET proteins because the pre-MET embryo is transcriptionally incompetent. This is evidenced by the fact that the nuclear transfer embryos progress through the MET and that the first readily observable

-14-

signs of differentiation, compaction and blastocoel formation, occur as in normal embryos.

Furthermore, the bovine recipient oocyte is apparently capable of supporting nuclear remodeling in the donor cells of various species. Nuclear remodeling refers to the series of changes that occur to the nucleus of the donor cell after nuclear transfer. (For review see Fulka et al., 1996; Prather and First, 1990.) The morphological and biochemical changes in the nucleus are probably due to factors present in the cytoplasm of the recipient oocyte. These changes include the disappearance of differentiated nucleoli, swelling of the transferred nucleus, and in amphibians the exchange of proteins between the nucleus and cytoplasm. These changes are probably necessary to support further development. Again, the bovine recipient oocyte apparently has the necessary proteins to support this remodeling in other species.

Third, the bovine recipient oocyte is probably capable of supporting correct changes in methylation of the DNA of the donor cells of the various species. DNA methylation has been shown to be a mechanism for the regulation of gene expression in mammals. (For review see Brandeis et al., 1993.) The DNA of mouse gametes is heavily methylated. Upon fertilization the DNA of the newly formed zygote is demethylated. At gastrulation, a bimodal pattern of methylation is formed with CpG island containing genes being demethylated and most other genes being methylated. Tissue specific genes are demethylated as differentiation continues. The bovine recipient oocyte may be able to support the initial wave of demethylation as evidenced by the developmental competence of the nuclear transfer embryos. After the MET, the genome of the donor cell probably exerts itself to establish correct methylation patterns.

In the preferred embodiment, fusion of the recipient oocyte and donor nucleus is performed by



-15-

electric pulse. In the most preferred embodiment, the bovine recipient oocyte is fused to the bovine donor cells or donor cells of other species by a double electric pulse in sucrose or sorbitol fusion medium in a Zimmerman Electrofusion Instrument (CGA Corporation, Chicago, Illinois). The method is similar to that described in Prather et al., U.S. Patent 4,994,384. The Electrofusion Instrument may be adjusted as follows:

10

Fusion voltage: 90 volts (DC, 1.8 KV/cm);

Electrode distance: 0.5 millimeter;

Alignment voltage: 1-5 volts (AC);

Alignment frequency: 1000 KHZ;

15

Pulse duration: 30 microseconds, two pulses;

Postfusion alignment time: 5 seconds

Sorbitol fusion medium consists of 0.25 M d-sorbitol in medical grade H<sub>2</sub>O (Baxter), 100  $\mu$ M Ca-acetate, 0.5 mM Mg-acetate, and 1 g/l Fatty Acid Free BSA (pH 7.2 at room temperature, 255 mOsm). Sucrose fusion medium consists of 95.84 g/l sucrose, 0.107 g/L Mg-acetate-4-H<sub>2</sub>O, 0.016  $\mu$ M Ca-acetate, 0.174 g/l KH<sub>2</sub>PO<sub>4</sub> anhydrous, 0.031 g/l glutathione, and 0.01 g/l Fatty Acid Free BSA, ELISA grade (260-290 mOsm, pH 7.0). Other fusion media may be used such as Zimmerman Cell Fusion Medium, mannitol, and phosphate buffer solution.

25

Alternatively, fusion may be accomplished by treatment of the donor cell and recipient oocyte with polyethylene glycol as described in Sims and First, 1994. Fusion may be accomplished also by use of Sendai virus. (Graham, 1969.)

30

The next step in cloning by nuclear transfer is activation of the nuclear transfer embryo. "Nuclear transfer embryo" means the embryo which results from the fusion of the donor cell and recipient oocyte. Normally, the oocyte is activated by the sperm when fertilized. The interaction of the sperm with the

35

-16-

oocyte cell membrane causes an elevation of intracellular calcium which starts a cascade of events including the release of the new zygote from Metaphase II arrest. Activation by fertilization is followed by pronuclear formation. Since nuclear transfer embryos are not fertilized, an activation stimulus is needed to release the nuclear transfer embryo from Metaphase II arrest. In traditional nuclear transfer protocols, the electric pulse used to fuse the donor cell and recipient oocyte acts as the activation stimulus. (Willadsen, 1986; Robl et al., 1986; Prather et al., 1987; Campbell et al., 1996.) It is necessary that sheep recipient oocytes and bovine recipient oocytes be aged in order for an electric pulse to be sufficient to activate the resulting nuclear transfer embryo. (Prather et al., 1987; Campbell et al., 1996.)

In a preferred embodiment, the nuclear transfer embryo is activated by a stimulus sufficient to release the nuclear transfer embryo from Metaphase II arrest. "Activate", as used herein, means the use of a stimulus sufficient to release the nuclear transfer embryo from Metaphase II arrest. In an alternative embodiment, a bovine nuclear transfer embryo is activated by the fusion electric pulse described above at 35 to 52 hours after the beginning of in vitro culture.

Recently, a method has been described for activating bovine oocytes as early as 10 hours after the beginning of in vitro culture. (Susko-Parish et al., U.S. Patent 5,496,720, incorporated herein by reference.) This method comprises the sequential incubation of an oocyte or nuclear transfer embryo with a calcium ionophore such as ionomycin, and a serine-threonine kinase inhibitor such as 6-dimethylaminopurine (DMAP). Treatment with ionomycin causes an increase in intracellular calcium similar to that normally caused by the sperm at fertilization. The role of DMAP is less clear, but it may act by inhibiting the serine-threonine kinase c-Mos which is

-17-

thought to stabilize the activity of p34cdc2 which is activated by phosphorylation on serine and threonine residues and inactivated by phosphorylation on tyrosine residues. High p34cdc2 activity is generally associated with Metaphase II arrest. Since the activity of c-Mos is inhibited, p34cdc2 activity is down regulated via the absence of an entity able to maintain the activating phosphorylations on p34cdc2's serine and threonine residues. One reason that aged oocytes may be activated solely by electrical pulse is that p34cdc2 activity in these cells has declined. All that is needed to activate aged oocytes is the elevation of intracellular calcium by one of several methods. To overcome the high levels of p34cdc2 activity in younger oocytes, both elevation of intracellular calcium and inhibition of serine-threonine kinases appear to be required.

In the preferred embodiment, nuclear transfer embryos are activated at an appropriate time (preferably about 4 to 16 hours after fusion) by increasing the concentration of intracellular calcium and incubation with a serine-threonine kinase inhibitor. In the most preferred embodiment, nuclear transfer embryos comprised of fused bovine recipient oocytes and bovine donor cells or donor cells from other species are activated by sequential incubation with ionomycin and DMAP at about 28 hours after the beginning of in vitro culture. Briefly, the nuclear transfer embryos are exposed to 5  $\mu$ M ionomycin (5 mM stock in DMSO; Calbiochem, La Jolla, CA) in TL-HEPES (1 mg/ml fatty acid free BSA) for 4 minutes. The nuclear transfer embryos are then diluted into TL-HEPES (30 mg/ml fatty acid free BSA) for five minutes and then further diluted into TL-HEPES (1 mg/ml Fraction V BSA). The nuclear transfer embryos are then moved to 50  $\mu$ l drops under paraffin oil of CRLaa with 1.9 mM DMAP and incubated for 4 hours at 39° C, 5% CO<sub>2</sub> in air, high humidity. After this incubation, embryos are diluted

-18-

in TL HEPES with 1 mg/ml BSA and then transferred into CRlaa media and cultured at 39° C, 5% CO<sub>2</sub> in air, high humidity. Other methods which may be used to increase intracellular calcium include ethanol treatment, A23187, electric shock and caged chelators. Other serine-threonine kinase inhibitors which may be used include staurosporine, 2-aminopurine, and sphingosine. For a further description of these methods see Susko-Parrish et al., U.S. Patent 5,496,720, incorporated herein by reference.

There are two main advantages to the use of the ionomycin/DMAP activation regimen. First, the nuclear transfer embryo may be activated at the time when developmental competence is highest as demonstrated by studies of the fertilization of bovine oocytes. Traditional cloning techniques require that the recipient oocyte be aged before fusion and activation. The use of aged oocytes may compromise the development of the resulting nuclear transfer embryo. Second, p34cdc2 has been shown to have a positive effect on nuclear remodeling. (Fulka et al., 1996.) The nuclei of nuclear transfer embryos which are exposed to a cellular environment with high levels of p34cdc2 activity remodel more completely and more quickly than when fusion occurs to recipients that have already been preactivated or which have low levels of p34cdc2 activity. One reason that aged oocytes may be activated solely by electrical pulse is that p34cdc2 activity in these cells has declined. Since p34cdc2 activity has declined in aged recipient oocytes, nuclear remodeling may not be complete. With the ionomycin/DMAP regimen, at the time of fusion, which occurs prior to activation, the donor cell nucleus is exposed to high levels of p34cdc2 activity which is beneficial to nuclear remodeling.

Temporal separation of the fusion and activation steps allows the nuclear transfer embryo sufficient time to remodel its nucleus. In the most preferred

-19-

embodiment, fusion occurs at 16 hours after the beginning of in vitro culture while activation occurs 28 hours after the beginning of in vitro culture. This procedure differs from previously described nuclear transfer techniques where fusion and activation are simultaneous. The examples verify that this method results in an approximately 2 fold increase to blastocyst than previously described somatic cell cloning methods.

After activation of the nuclear transfer embryo, it is necessary to either culture the embryo in vitro or transfer the embryo to an appropriate recipient animal. In the preferred embodiment, nuclear transfer embryos comprised of bovine recipient oocytes and bovine donor cells are cultured until the blastocyst stage in vitro and then transferred to a recipient. In vitro culture is carried out as described in Sims and First, 1994 and Rosenkrans et al., 1993. Briefly, nuclear transfer embryos may be cultured in a defined medium, CR1aa or CR2 for 7-8 days at 39° C in 5% CO<sub>2</sub>/95% air in high humidity. CR1aa is comprised of 5 mM hemicalcium lactate, 114.7 mM sodium chloride, 3.1 mM potassium chloride, 26.2 mM sodium bicarbonate, and 3 mg/ml fatty acid free bovine serum albumin. (Rosenkrans et al., 1993.)

#### EXAMPLES

##### Methods Common to all Experiments

Oocyte maturation- In vitro matured bovine oocytes may be prepared by the methods of Sirard et al. (1988), Parrish et al. (1988) and Rosenkrans and First (1993). Briefly, ovaries are collected at a slaughterhouse and transported to the laboratory in normal saline at 30° to 34° C. Transport time varies from 2 to 6 hours. Oocytes from small follicles of 1 to 6 mm in diameter are aspirated. The immature oocytes are then washed

-20-

three times in TL-HEPES without glucose and to which 0.22 mM pyruvate and 1 mg/ml BSA have been added. The immature oocytes are then placed in 50  $\mu$ l drops of maturation medium under paraffin oil. Maturation  
5 medium is composed of TC199 containing 10 percent heat treated fetal calf serum, 0.22 mM pyruvate, 0.5  $\mu$ g/ml FSH-P (Scherring-Plough Animal Health Corp. Kenilworth, N.J.) and 1  $\mu$ g/ml estradiol. Ten oocytes are placed in each drop and incubated overnight at 39° C, in high  
10 humidity atmosphere comprised of 5% CO<sub>2</sub> and 95% air.

Micromanipulation- Enucleation was performed at 16-20 hours after the beginning of in vitro culture. The procedure for enucleation is as follows and is  
15 described in Prather et al., U.S. Patent 4,994,384, incorporated herein by reference. Cumulus cells are removed from the oocytes by placing the oocytes in TL-HEPES medium supplemented with 2.0 milligrams per milliliter of hyaluronidase and then gently aspirating  
20 the oocytes with a fire-polished pipette tip. (Sims and First, 1994.) Oocytes to be enucleated are then placed in culture dishes in 100 microliter drops under oil of TL-HEPES medium supplemented with calcium and magnesium. (See generally Sims and First, 1994.)  
25 Micromanipulation may be performed with a Nikon Diaphot microscope equipped Hoffman optics and Narishige micromanipulators. The recipient oocyte is held with a holding pipet with a 120  $\mu$ m outer diameter and 30  $\mu$ m inner diameter. Enucleation is performed by using a  
30 beveled, sharpened enucleation pipet with an outer diameter of approximately 25  $\mu$ m. The inner diameter may be varied according to the size of the donor cell which is used. To enucleate the oocyte, the enucleation pipet is inserted into the oocyte and a  
35 small amount of cytoplasm is removed just below the first polar body. Preferably, enucleation can be verified by staining the mature recipient oocyte with Hoechst 33342, visualizing DNA by excitation emission

-21-

upon exposure to ultraviolet light for a short time (i.e. 5 seconds), enucleating the recipient oocyte, and washing the oocyte 2 times in TL-HEPES. The donor cell is placed under the zona pellucidae of the recipient oocyte by use of the enucleation pipet.

Fusion- Fusion of the bovine, non-human primate, rat, sheep and porcine donor cells to the bovine recipient oocytes was performed at 16-24 hours after the beginning of in vitro culture. The bovine recipient oocyte is fused to the bovine donor cells or donor cells of other species by a double electric Zimmerman Electrofusion Instrument (CGA Corporation, Chicago, Illinois). The method is similar to that described in Prather et al., U.S. Patent 4,994,384. The Electrofusion Instrument was adjusted as follows:

Fusion voltage: 90 volts (DC, 1.8 KV/cm);  
Electrode distance: 0.5 millimeter;  
Alignment voltage: 1-5 volts (AC);  
Alignment frequency: 1000 KHZ;  
Pulse duration: 30  $\mu$ seconds, two pulses;  
Postfusion alignment time: 5 seconds

Two types of fusion media were utilized. The first consisted of 0.25 M d-sorbitol in medical grade H<sub>2</sub>O (Baxter), 100  $\mu$ M Ca-acetate, 0.5 mM Mg-acetate, and 1 g/l Fatty Acid Free BSA (pH 7.2 at room temperature, 255 mOsm). The second consisted of 95.84 g/l sucrose, 0.107 g/L Mg-acetate-4-H<sub>2</sub>O, 0.016  $\mu$ M Ca-acetate, 0.174 gA KH<sub>2</sub>PO<sub>4</sub> anhydrous, 0.031 g/l glutathione, and 0.01 g/l Fatty Acid Free BSA, ELISA grade (260-290 mOsm, pH 7.0).

Cell culture- Somatic cells were obtained from bovine, sheep, non-human primate (Rhesus), porcine and rat ear skin samples, removed from live animals. The skin samples were placed into PBS (phosphate buffered

-22-

saline, pH ) supplemented with antibiotic/antimycotic (Gibco), and washed 3 times. All hair was manually removed and the sample cut into small pieces. The cut samples were then placed in 0.05% trypsin/1 mM EDTA solution (Gibco) for 40 minutes at 30° C and stirred frequently during incubation. The samples were then centrifuged (10 minutes, 1000 rpm) and supernatant containing single cells removed to remove undigested pieces of tissue. The final supernatant was placed into alpha-MEM (Gibco), supplemented 10% fetal calf serum (HiClone) and antibiotic/antimycotic (Gibco) and cultured in tissue flasks (Falcon, 25 cm<sup>2</sup>) at 37° C for at least a week. After a monolayer of cells had formed, the medium was removed and the monolayer washed with Ca-Mg free PBS once and replaced by trypsin/EDTA solution and incubated at 37° C for 2-3 minutes. Cells are then desegregated by pipetting and centrifuged as described above. The pellet of cells was diluted 1:2 with alpha-MEM and placed into two flasks. All subsequent passages were performed in the same manner every 2 days thereafter.

For nuclear transfer, the cells were removed from flasks as described for passaging and placed into a 35 mm Petri dish (Falcon tissue culture) and grown in alpha-MEM for 2-12 days in the absence of any serum. On the day of nuclear transfer, the cells grown in the absence of sera were trypsinized as described above, washed in Ca-Mg free PBS and placed into TL HEPES supplemented with 0.1 -3 mg/ml BSA Fraction V, 0.2 mM Na-pyruvate and 25 µg/ml gentamicin.

Activation of nuclear transfer embryos- Activation was performed at 24-28 hours after the beginning of in vitro culture. The nuclear transfer embryos were exposed to 5 µM ionomycin (5 mM stock in DMSO; Calbiochem, La Jolla, CA) in TL-HEPES (1 mg/ml fatty acid free BSA) for 4 minutes. The nuclear transfer embryos were then diluted into TL-HEPES (30 mg/ml fatty



-23-

acid free BSA) for five minutes and then twice diluted into TL-HEPES (3 mg/ml Fraction V BSA). The nuclear transfer embryos were then moved to 50 microliter drops under paraffin oil of CR1aa supplemented with 1.9 mM DMAP and incubated for 4 hours at 39° C, 5% CO<sub>2</sub> in air, high humidity.

Culture after activation- After removal from DMAP, nuclear transfer embryos were diluted twice in TL HEPES supplemented with 3 mg/ml BSA Fraction V. All embryos, regardless of species, were cultured in CR1aa. CR1aa is comprised of 5 mM hemicalcium lactate, 114.7 mM sodium chloride, 3.1 mM potassium chloride, 26.2 mM sodium bicarbonate, and 3 mg/ml fatty acid free bovine serum albumin. (Rosenkrans et al., 1993.) Incubations were conducted at 39° C, 5% CO<sub>2</sub> in air, high humidity. After three days of culture, 10% heat treated FCS was added to all embryos, regardless of species and the embryos allowed to develop to blastocyst.

20

### Results

Nuclear transfer embryos were constructed from bovine recipient oocytes and the following cell lines: bovine adult fibroblasts (BAF), bovine fetal fibroblasts (BFF), sheep adult fibroblasts (SAF), porcine adult fibroblasts (PAF), non-human primate adult fibroblasts (MAF) and rat adult fibroblasts (RAF). The results are summarized in Table 1.

30

- 24 -

Table 1: Ability of donor cells to initiate embryonic development after nuclear transfer into a recipient bovine oocyte.

Donor Cell Line	Nuclear Transfer (N)	Fused (N)	Lysed or not fused (N)	CLV.* (%/fused)	CM+BL** (%/C.V.)
BAF	220	145	75	81/145	22/81
BFF	82	51	31	31/51	18/31
SAF	327	182	135	120/135	38/120
PAF	93	36	57	32/57	12/32
RAF	108	33	75	37/	na
MAF	30	28	2	24/28	14/24

\* CLV. - number of embryos undergoing cleavage

\*\* CM-BL- number of embryos developing to compact morula or blastocyst

Column 1 indicates the donor cell line, column 2, Nuclear Transfer, indicates the number of nuclear transfers attempted, column 3, Fused, indicates the number of fused nuclear transfer embryos produced, column 4, Lysed, indicates the number of recipient oocytes which lysed or where fusion did not occur, column 5, CLV, indicates the number nuclear transfer embryos which cleaved and is expressed as a percentage of fused nuclear transfer embryos, column 6, CM+BL, indicates the number of nuclear transfer embryos developing to compacted morula or blastocyst and is expressed as a percentage of cleaved embryos.

These results indicate the bovine oocyte can be utilized as the recipient oocyte in combination differentiated donor cells from cattle, pigs, monkeys, rats and sheep to create nuclear transfer embryos. The methods employed lead to a greater efficiency of development to compacted morula and blastocyst than previously described somatic cell cloning procedures. Wilmut et al., reported the development of 29 embryos

-25-

to compacted morula or blastocyst out of 277 fused nuclear transfer embryos (11.7%) when using somatic sheep cells. Here the researchers report the development of 38 compacted morulae or blastocysts out of 182 fused nuclear transfer embryos (20.9%). An increase in efficiency of about two fold.

To date, 13 embryos derived from bovine adult fibroblasts have been transferred to recipient cows, 8 embryos derived from sheep adult fibroblasts have been transferred to sheep, and 20 embryos derived from rat adult fibroblasts have been transferred to rats.

#### REFERENCES

- 15 Anderson, *Animal Biotechnology* 3, 165-175 (1992).  
Barnes and First, Embryonic Transcription in in vitro Cultured Covine Embryos, *Molecular Reproduction and Development* 29, 117-123 (1991).  
20 Brandeis et al., Dynamics of DNA Methylation During Development, *BioEssays*, 15 709-713 (1993)  
Campbell et al., Sheep Cloned by Nuclear Transfer from a Cultured Cell Line, *Nature*, 380 64-66 (1996)  
Capecchi, Altering the Genome by Homologous Recombination, *Science* 244, 1288-1292 (1989).  
25 Connover et al., Stage Specific Expression of a Family of Proteins that are Major Products of Gene Activation in the Mouse Embryo, *Developmental Biology* 147, 403-414 (1991).  
Crosby et al., Control of Protein Synthesis During Early Cleavage of Sheep Embryos, *Journal of Reproduction and Fertility* 82, 769-775 (1988).  
30 DiBerardino, Genornic Potential of Differentiated Cells Analyzed by Nuclear Transplantation *American Zoology*, 27, 623-644 (1987).  
35 Dominko and First, Timing of Meiotic Progression in Bovine Oocytes and Its Effect on Early Embryo Development, *Molecular Reproduction and Development*, 47, 456-467 (1997).

-26-

- First et al., Systems for Production of Calves from Cultured Bovine Embryonic Cells, *Reproduction, Fertility and Development*, 6 553-562 (1994).
- 5 Fulka et al., Nuclear Transplantation in Mammals: Remodeling of Transplanted Nuclei Under the Influence of Maturation Promoting Factor, *BioEssays*, 18 835-840 (1996).
- Graham, The Fusion of Cells with One and Two Cell Mouse Embryos, *Wistar. Inot. Symp. Monogr.*, 9, 19
- 10 (1969).
- Gurdon, Nuclear Transplantation in Eggs and Oocytes, *Journal of Cell Science* 4 (Supp.) 287-318 (1987)
- Joyner, Gene Targeting and Gene Trap Screens Using Embryonic Stem Cells: New approaches to Mammalian
- 15 Development, *BioEssays* 13, 649-658 (1991).
- Koller et al., Proceedings of the National Academy of Sciences, USA 86, 8927-8931 (1989).
- McGrath and Solter, Nuclear Transplantation in the Mouse Embryo by Microsurgery and Cell Fusion,
- 20 *Science* 220, 1300-1302 (1983).
- Meng et al., Full Term Development of Rhesus Monkeys Produced by Nuclear Transfer, *Biology of Reproduction* (Supp.) 56, 87 (1997).
- Parrish et al., Capacitation of Bovine Sperm by
- 25 Heparin, *Biology of Reproduction* 38, 1171-1180 (1988)
- Prather et al., Nuclear Transplantation in Early Pig Embryos, *Biology of Reproduction*, 41 414 418 (1989).
- 30 Prather et al., Nuclear Lamin Antigens: Porcine and Bovine Early Developmental Regulation, *Biology of Reproduction*, 41 414-418 (1989).
- Prather and First, Cloning Embryos by Nuclear Transfer, *Journal of Reproduction and Fertility*, Supplement
- 35 41, 125-134 (1990).
- Robl et al., Nuclear Transplantation in Bovine Embryos, *Journal of Animal Science*, 64 642-647 (1986)
- Robl et al., Nuclear Transplantation in the Bovine

-27-

- Embryo: Assessment of Donor Nuclei and Recipient Oocyte, *Biology of Reproduction*, 37 859-866 (1987).
- 5       Rosenkrans et al., *Journal of Animal Science* 72, 634-637 (1993).
- Sims and First, Production of Calves by Transfer of Nuclei from Cultured Inner Cell Mass Cells, *Proceedings of the National Academy of Sciences USA*, 90 6143-6147 (1993).
- 10       Sirard et al., The Culture of Bovine Oocytes to Obtain Developmentally Competent Embryos, *Biology of Reproduction* 39, 546-552 (1988).
- Stanton et al., Germ Line Transmission of an Inactive N-myc Allele Generated by Homologous Recombination
- 15       in Mouse Embryo Stem Cells, *Molecular and Cellular Biology* 10, 6755-6758 (1990)
- Stewart, in *Animal applications of Research in Mammalian Development*, eds. Pederson, R. A., McLaren, A., and First, N. (Cold Spring Harbor
- 20       Laboratory Press, Plainview, NY) 267-283 (1991).
- Stice and Robl, Nuclear Reprogramming in Nuclear Transplant Rabbit Embryos, *Biology of Reproduction* 39, 657-664 (1988).
- 25       Susko-Parish et al., Inhibition of Protein Kinases after an Induced Calcium Transient Causes Transition of Bovine Oocytes to Embryonic Cycles without Meiotic Completion, *Developmental Biology* 166, 729-739 (1994).
- 30       Telford et al., Transition from Maternal to Embryonic Control in Early Mammalian Development: A Comparison of Species, *Molecular Reproduction and Development* 26, 90-100 (1990).
- Willadsen, Nuclear Transplantation in Sheep Embryos,
- 35       *Nature*, 320 63-65 (1986).
- Wilmut et al, Viable Offspring Derived from Fetal and Adult Mammalian Cells, *Nature*, 385 810813 (1997).

-28-

CLAIMS

What is claimed is:

- 5  
1. A method of producing nuclear transfer bovine embryos from bovine donor cells and bovine recipient oocytes comprising:  
inducing bovine donor cells to undergo G<sub>0</sub> arrest;  
10 fusing said bovine donor cell to an enucleated bovine recipient oocyte to create a nuclear transfer embryo;  
and activating said nuclear transfer embryo.
- 15 2. The method of claim 1 wherein said G<sub>0</sub> arrest of donor cells is induced by culture in low serum medium;
- 20 3. The method of claim 1 wherein said bovine donor cells are selected from the group consisting of embryonic derived cells, germ cells, somatic cells, and genetically modified cells.
- 25 4. The method of claim 1 wherein said enucleated bovine recipient oocyte is selected from the group of bovine oocytes undergoing nuclear maturation within 16 hours of beginning in vitro culture.
- 30 5. The method of claim 1 wherein said enucleated bovine recipient oocyte and said bovine donor cell are fused by electric pulse to form a nuclear transfer embryo.
- 35 6. The method of claim 1 wherein said fusion is performed 16-24 hours after the beginning of in vitro culture.

-29-

7. The method of claim 1 wherein said nuclear transfer embryo is activated by elevating intracellular calcium and then incubating with a serine threonine kinase inhibitor.
- 5 8. The method of claim 7 wherein intracellular calcium is elevated by incubation with ionomycin and the serine threonine kinase inhibitor is DMAP.
- 10 9. The method of claim 1 wherein said activation is 16 to 32 hours after the beginning of in vitro culture.
10. The method of claim 5 wherein said fusion is 16-52 hours after the beginning of in vitro culture.
- 15 11. An embryo produced by the method of claim 1.
12. A method of producing nuclear transfer embryos from donor cells of one species and recipient oocytes from another species comprising:
- 20 inducing the donor cells to undergo G<sub>0</sub> arrest;  
fusing said donor cell to an enucleated recipient oocyte of another species to create a nuclear transfer embryo;
- 25 and activating said nuclear transfer embryo.
13. The method of claim 12 wherein said G<sub>0</sub> arrest of donor cells is induced by culture in low serum medium.
- 30 14. The method of claim 12 wherein said donor cells are selected from the group consisting of embryonic derived cells, germ cells, somatic cells, and genetically modified cells.
- 35 15. The method of claim 12 wherein said enucleated recipient oocyte is an enucleated bovine recipient oocyte.

-30-

16. The method of claim 15 wherein said enucleated bovine recipient oocyte is selected from the group of bovine oocytes undergoing nuclear maturation within 16 hours of beginning in vitro culture.

5

17. The method of claim 12 wherein said enucleated bovine recipient oocyte and said donor cell are fused by electric pulse to form a nuclear transfer embryo.

10

18. The method of claim 12 wherein said fusion is performed 16-32 hours after the beginning of in vitro culture.

15

19. The method of claim 12 wherein said nuclear transfer embryo is activated by elevating intracellular calcium and the incubating with a serine threonine kinase inhibitor.

20

20. The method of claim 19 wherein intracellular calcium is elevated by incubation with ionomycin and the serine threonine kinase inhibitor is DMAP.

25

21. The method of claim 12 wherein said activation is 16-32 hours after the beginning of in vitro culture.

22. The method of claim 17 wherein said fusion is 16-52 hours after the beginning of in vitro culture.

30

23. An embryo produced by the method of claim 12.

24. A method for producing nuclear transfer bovine embryos from bovine donor cells and bovine recipient oocytes comprising:

35

culturing bovine donor cells selected from the group consisting of embryonic derived cells, somatic cells, germ cells, and genetically modified cells in low serum medium so that said donor cells are induced to arrest in the G<sub>0</sub> stage of the cell cycle;



-31-

selecting a bovine recipient oocyte which has completed nuclear maturation before 16 hours from the beginning of in vitro culture;

5 enucleating said bovine recipient oocyte after 16-32 hours of in vitro culture;

placing said donor cell under the zona pellucida of said enucleated oocyte so that said donor cell contacts said enucleated oocyte.

10 fusing said donor cell with said enucleated oocyte by electric pulse at 16-32 hours after the beginning of in vitro culture to create a nuclear transfer embryo;

and activating said nuclear transfer embryo by sequential incubation with ionomycin and 6-dimethylaminopurine at 16 to 32 hours after the beginning of in vitro culture.

25. The embryo produced by the process of claim 24.

26. A method of producing nuclear transfer embryos from a donor cell of one species and a bovine recipient oocyte comprising:

20 culturing non-bovine donor cells selected from the group consisting of embryonic derived cells, somatic cells, germ cells, and genetically modified cells in low serum medium so that said donor cells are induced to arrest in the G<sub>0</sub> stage of the cell cycle;

selecting a bovine recipient oocyte which has completed nuclear maturation before 16 hours from the beginning of in vitro culture;

30 enucleating said bovine recipient oocyte after 16-32 hours of in vitro culture;

placing said donor cell under the zone pellucida of said enucleated oocyte so that said donor cell contacts said enucleated oocyte;

35 fusing said donor cell with said enucleated oocyte by electric pulse at 16-32 hours after the beginning of in vitro culture to create a nuclear transfer embryo;

and activating said nuclear transfer embryo by

-32-

sequential incubation with ionomycin and  
6-dimethylaminopurine at 16 to 32 hours after the  
beginning of in vitro culture.

5        27. The embryo produced by the process of claim 25.

28. A nuclear transfer embryo comprising  
cytoplasm derived from a bovine oocyte, bovine  
cytoplasm derived from a differentiated bovine cell,  
10       cell membrane derived from a bovine oocyte, cell  
membrane derived from a differentiated bovine cell and  
nuclei derived from a differentiated bovine cell.

29. A nuclear transfer embryo comprising cytoplasm and  
15       cell membrane from one species and differentiated  
cytoplasm, differentiated cell membrane, and nuclei  
derived from a differentiated cell of another species.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/00, A01K 67/027</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/05266</b> <b>(43) International Publication Date:</b> <b>4 February 1999 (04.02.99)</b>
<b>(21) International Application Number:</b> PCT/US98/15387 <b>(22) International Filing Date:</b> 24 July 1998 (24.07.98) <b>(30) Priority Data:</b> 60/053,103 26 July 1997 (26.07.97) US <b>(71) Applicant (for all designated States except US):</b> WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, P.O. Box 7365, Madison, WI 53707-7365 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FIRST, Neal, L. [US/US]; 3157 Highway 78, Mt. Horeb, WI 53572 (US). DOMINKO, Tanja [SL/US]; 2320 White Oak Trail, Oregon, WI 53575 (US). MITALIPOUA, Maissam [US/US]; 2130 University Avenue #6, Madison, WI 53705 (US). <b>(74) Agent:</b> BARTA, Kent, S.; Michael Best & Friedrich LLP, 100 East Wisconsin Avenue, Milwaukee, WI 53202-4108 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW. ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>With amended claims and statement.</i> <b>(88) Date of publication of the international search report:</b> 15 April 1999 (15.04.99)  <b>Date of publication of the amended claims and statement:</b> 23 September 1999 (23.09.99)	
<b>(54) Title:</b> TRANS-SPECIES NUCLEAR TRANSFER  <b>(57) Abstract</b>  A method of producing cloned nuclear transfer embryos from differentiated donor cells is described. The method includes culturing differentiated donor cells in low serum medium and using bovine oocytes as recipient oocytes for donor cells from different species. This method may be used to produce genetically identical animals and transgenic animals.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## AMENDED CLAIMS

[received by the International Bureau on 23 April 1999 (23.04.99);  
original claims 1-29 replaced by amended claims 1-15 (3 pages)]

1. A method of producing nuclear transfer embryos from donor cells of one species and recipient oocytes from another species comprising:
  - inducing the donor cells to undergo G<sub>0</sub> arrest;
  - fusing said donor cell to an enucleated recipient oocyte of another species to create a nuclear transfer embryo;
  - and activating said nuclear transfer embryo.
2. The method of claim 1 wherein said G<sub>0</sub> arrest of donor cells is induced by culture in low serum medium.
3. The method of claim 1 wherein said donor cells are selected from the group consisting of embryonic derived cells, germ cells, somatic cells, and genetically modified cells.
4. The method of claim 1 wherein said enucleated recipient oocyte is an enucleated bovine recipient oocyte.
5. The method of claim 4 wherein said enucleated bovine recipient oocyte is selected from the group of bovine oocytes undergoing nuclear maturation within 16 hours of beginning in vitro culture.
6. The method of claim 1 wherein said enucleated bovine recipient oocyte and said donor cell are fused by electric pulse to form a nuclear transfer embryo.
7. The method of claim 1 wherein said fusion is performed 16-32 hours after the beginning of in vitro culture.
8. The method of claim 1 wherein said nuclear transfer embryo is activated by elevating intracellular calcium and the incubating with a serine threonine kinase inhibitor.

9. The method of claim 8 wherein intracellular calcium is elevated by incubation with ionomycin and the serine threonine kinase inhibitor is DMAP.
10. The method of claim 1 wherein said activation is 16-32 hours after the beginning of in vitro culture.
11. The method of claim 6 wherein said fusion is 16-52 hours after the beginning of in vitro culture.
12. An embryo produced by the method of claim 1.
13. A method of producing nuclear transfer embryos from a donor cell of one species and a bovine recipient oocyte comprising:
- culturing non-bovine donor cells selected from the group consisting of embryonic derived cells, somatic cells, germ cells, and genetically modified cells in low serum medium so that said donor cells are induced to arrest in the G<sub>0</sub> stage of the cell cycle;
  - selecting a bovine recipient oocyte which has completed nuclear maturation before 16 hours from the beginning of in vitro culture;
  - enucleating said bovine recipient oocyte after 16-32 hours of in vitro culture;
  - placing said donor cell under the zone pellucida of said enucleated oocyte so that said donor cell contacts said enucleated oocyte;
  - fusing said donor cell with said enucleated oocyte by electric pulse at 16-32 hours after the beginning of in vitro culture to create a nuclear transfer embryo;
  - and activating said nuclear transfer embryo by sequential incubation with ionomycin and 6-dimethylaminopurine at 16 to 32 hours after the beginning of in vitro culture.
14. The embryo produced by the process of claim 13.

15. A nuclear transfer embryo comprising cytoplasm and cell membrane from one species and differentiated cytoplasm, differentiated cell membrane, and nuclei derived from a differentiated cell of another species.

**STATEMENT UNDER ARTICLE 19**

The original claims 1-29 have been amended, by canceling original claims 1-11, 24, 25, and the remaining original claims 12-23, 26, 27 and 29 have been renumbered as claims 1-15. The claims after amendment are now directed solely to a method for producing nuclear transfer embryos from donor cells of one species and recipient oocytes from another species.



## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/US 98/15387

 A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C12N15/00 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)  
 IPC 6 A01K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 07669 A (ROSLIN INST EDINBURGH ; CAMPBELL KEITH HENRY STOCKMAN (GB); WILMUT) 6 March 1997 cited in the application	1-3, 5, 9, 11, 28
Y	see page 2, line 26 - page 5, line 24 see page 7, line 13 - line 24; examples 1, 4 see page 8, line 24 - page 9, line 12; claims	1-29
Y	STICE S L ET AL: "PLURIPOTENT BOVINE EMBRYONIC CELL LINES DIRECT EMBRYONIC DEVELOPMENT FOLLOWING NUCLEAR TRANSFER" BIOLOGY OF REPRODUCTION, vol. 54, no. 1, January 1996, pages 100-110, XP000671718 cited in the application see the whole document	1-3, 5-11, 24, 25, 28

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

4 February 1999

Date of mailing of the international search report

23 02 99

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Chambonnet, F

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/15387

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SIMS M ET FIRST: "PRODUCTION OF CALVES BY TRANSFER OF NUCLEI FROM CULTURED INNER CELL MASS CELLS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, no. 13, June 1993, pages 6143-6147, XP000604993 cited in the application see the whole document ---	1-3,5-7, 10
Y	CHESNE P ET AL: "NUCLEAR TRANSFER IN CATTLE: BIRTH OF CLONED CALVES AND ESTIMATION OF BLASTOMERE TOTIPOTENCY IN MORULAE USED AS A SOURCE OF NUCLEI" LIFE SCIENCES, vol. 316, 1993, pages 487-491, XP000197855 see the whole document ---	1-3
Y	US 5 496 720 A (SUSKO-PARRISH JOAN L ET AL) 5 March 1996 cited in the application see the whole document ---	4-10
P,Y	WO 98 30683 A (UNIV MASSACHUSETTS A PUBLIC IN) 16 July 1998 see the whole document ---	3-11
A	WOLFE, B.A. ET AL.: "Embryos produced by the transfer of caprine nuclei to enucleated bovine oocytes are capable of cleavage but not to develop to blastocysts" BIOLOGY OF REPRODUCTION, vol. 50, no. suppl. 1, 1994 - 1972, page 72 XP002092259 see abstract 72 ---	12
A	HEYMAN Y ET AL: "CLONING OF DOMESTIC SPECIES" ANIMAL REPRODUCTION SCIENCE, vol. 42, no. 1/04, 1996, pages 427-436, XP000671884 see page 429, paragraph 2.5 ---	12
P,Y	WO 98 07841 A (UNIV MASSACHUSETTS) 26 February 1998 see the whole document ---	12-23, 26,27,29
T	ZAWADA, W.M. ET AL.: "Somatic cell cloned transgenic bovine neurons for transplantation in Parkinsonian rats" NATURE MEDICINE, vol. 4, no. 5, May 1998, pages 569-574, XP002092260 see the whole document -----	1

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 98/15387

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11 24 25 28

a method for producing nuclear transfer bovine embryos from bovine donor cells and bovine recipient oocytes comprising:  
inducing bovine donor cells to undergo G0 arrest;  
fusing said bovine donor cell to an enucleated bovine recipient oocyte to create a nuclear transfer embryo;  
and activating said nuclear transfer embryo; bovine embryo produced by this method; nuclear transfer bovine embryo comprising cytoplasm derived from a bovine oocyte, bovine cytoplasm derived  
from a differentiated bovine cell, cell membrane derived  
from a bovine oocyte, cell membrane derived  
from a differentiated bovine cell and nuclei derived  
from a differentiated bovine cell

2. Claims: 12-23 26 27 29

a method for producing nuclear transfer embryos from donor cells of one species and recipient oocytes from another species comprising:  
inducing the donor cells to undergo G0 arrest;  
fusing said donor cell to an enucleated recipient oocyte of another species to create a nuclear transfer embryo;  
and activating said nuclear transfer embryo; embryo produced by this method; nuclear transfer interspecies embryo comprising cytoplasm and cell membrane derived from one species and differentiated cytoplasm, differentiated cell membrane and nuclei derived from a differentiated cell from another species

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In. ational Application No

PCT/US 98/15387

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9707669	A	06-03-1997	AU 6831096 A	19-03-1997
			CA 2229568 A	06-03-1997
			CZ 9800608 A	15-07-1998
			EP 0849990 A	01-07-1998
			GB 2318578 A	29-04-1998
			NO 980845 A	29-04-1998
			PL 325331 A	20-07-1998
US 5496720	A	05-03-1996	US 5843754 A	01-12-1998
WO 9830683	A	16-07-1998	AU 6014598 A	03-08-1998
WO 9807841	A	26-02-1998	AU 4044397 A	06-03-1998